

The first of these is the fact that the majority of the cases of  
 this disease are reported from the West Coast of the United States,  
 particularly from California. It is also true that the disease is  
 more common in the summer months, and that it is more common  
 in the young than in the old. These facts are all in favor of the  
 theory that the disease is caused by a specific micro-organism,  
 and that it is transmitted from one person to another by direct  
 contact, or by the use of contaminated articles of personal  
 hygiene.

**Syphilis Venerea, seu Venereum**

This disease is characterized by the presence of a hard chancre  
 at the point of infection, and by the appearance of a rash  
 on the trunk and limbs. It is also characterized by the presence  
 of a fever, and by the enlargement of the lymphatic glands.  
 The disease is usually cured by the use of mercury, and by  
 the use of iodine. It is also cured by the use of arsenic,  
 and by the use of potassium iodide.

- (L. 1.) Syphilis Venerea
- (L. 2.) Syphilis Venerea
- (L. 3.) Syphilis Venerea
- (L. 4.) Syphilis Venerea
- (L. 5.) Syphilis Venerea

The disease is usually cured by the use of mercury, and by  
 the use of iodine. It is also cured by the use of arsenic,  
 and by the use of potassium iodide. The disease is usually  
 cured by the use of mercury, and by the use of iodine. It is  
 also cured by the use of arsenic, and by the use of potassium  
 iodide.

**Syphilis Venerea, seu Venereum**

This disease is characterized by the presence of a hard chancre  
 at the point of infection, and by the appearance of a rash  
 on the trunk and limbs. It is also characterized by the presence  
 of a fever, and by the enlargement of the lymphatic glands.  
 The disease is usually cured by the use of mercury, and by  
 the use of iodine. It is also cured by the use of arsenic,  
 and by the use of potassium iodide.